

The Problems of Time and Observables: Some Recent Mathematical Results

C. G. Torre [†]

Einstein's theory of gravitation can be viewed as a constrained Hamiltonian system [1]. The constraints are "first class", which indicates they generate "gauge transformations" on the phase space. In such systems it is important to separate cleanly the "pure gauge" content of the phase space from the "gauge invariant" information. In general relativity this leads, respectively, to the problem of time and the problem of observables. Progress on these problems represents an improvement in our understanding of the classical dynamics of the gravitational field, but these problems are particularly relevant to the construction of a quantum theory of gravity. The problem of time has been reviewed quite nicely by Kuchař [2]. The problem of observables has been extensively discussed in the literature, but it is hard to improve on the presentation due to Bergmann [3]. Both of these problems have been studied from a variety of viewpoints, but rigorous results have been obtained primarily in the context of simplified model systems. Here we give a couple of results on these problems in the context of the full vacuum theory.

The spacetime manifold M is assumed to have the topology $M = R \times \Sigma$, where Σ is a compact 3-manifold. We will consider the phase space Γ for gravitational dynamics to be a cotangent bundle over the space of Riemannian metrics on Σ . Dynamics takes place on the constraint surface $\bar{\Gamma}$ defined by the Hamiltonian and momentum constraints $\mathcal{H} = 0$ and $\mathcal{H}_a = 0$.

Our first result deals with the problem of time; specifically, we examine a proposal of Kuchař to solve the problem of time by identifying Γ with the phase space of some "parametrized field theory" [2], [4]. The idea is to isolate four fields X^α from Γ that can, at least on $\bar{\Gamma}$, be viewed as representing a spacelike embedding $X^\alpha: \Sigma \rightarrow M$. The constraints $\mathcal{H} = 0 = \mathcal{H}_a$ are then interpreted as conditions that identify the momenta P_α conjugate to the embedding with the energy-momentum densities of the remaining dynamical degrees of freedom. In order to investigate the viability of this scheme, we compare the constraint surface of a generic parametrized theory with $\bar{\Gamma}$. To this end, let $T^*\mathcal{E}$ denote a cotangent bundle over the space of embeddings $X^\alpha: \Sigma \rightarrow M$, and let Ω denote an infinite dimensional symplectic manifold. The phase space Ω is to represent the "true degrees of freedom" of the gravitational field; points of Ω will be labeled Z^A . The phase space Υ for a generic parametrized field theory is given by $\Upsilon = T^*\mathcal{E} \times \Omega$. Dynamics takes place on the constraint surface $\bar{\Upsilon}$, which is defined as follows. Let h_α denote four densities constructed from the embeddings X^α and dynamical variables Z^A . $\bar{\Upsilon}$ is then the subspace of Υ for which $P_\alpha + h_\alpha = 0$.

To implement the proposal of Kuchař we must find a canonical transformation (symplectic diffeomorphism) $\Phi: \Upsilon \rightarrow \Gamma$ such that $\Phi(\bar{\Upsilon}) = \bar{\Gamma}$. However, no such diffeomorphism exists [5]. The proof of this result rests on the fact that $\bar{\Upsilon}$ is a manifold, which can be seen

[†] Department of Physics, Utah State University, Logan, UT 84322-4415 USA

by applying the implicit function theorem. On the other hand, provided the topology of Σ does not prohibit the existence of vacuum spacetimes with Killing vectors, the constraint surface $\bar{\Gamma}$ is not a manifold [6]. Hence there can be no bijection identifying $\bar{\mathcal{T}}$ and $\bar{\Gamma}$. The essence of the difficulty with making this identification is that, in a spacetime with an isometry, it is impossible to distinguish embeddings that differ by an action of the isometry using only the intrinsic and extrinsic geometry of the embedded hypersurface. There are probably ways to mitigate this difficulty but, strictly speaking, general relativity is not a parametrized field theory.

Our second result concerns the existence of gravitational observables that can be expressed as spatial integrals of densities which are built locally from the canonical variables and their derivatives to any order. Recall that the constraint functions $(\mathcal{H}, \mathcal{H}_a)$, being “first class”, generate 1-parameter families of canonical transformations that leave $\bar{\Gamma}$ invariant. An observable is defined as a function $F: \Gamma \rightarrow \mathbb{R}$ whose restriction to $\bar{\Gamma}$ is invariant under the flow generated by the constraints. What this means in practice is that one looks for functions on Γ whose Poisson brackets with $\mathcal{H}$ and $\mathcal{H}_a$ vanish on $\bar{\Gamma}$. Any observable that vanishes on the constraint surface is called *trivial*. Any two observables that differ by a trivial observable will carry the same physical information, therefore we identify any two observables if their difference vanishes on $\bar{\Gamma}$. Next, recall that there are an infinite number of Hamiltonians that generate vacuum spacetimes from initial data on $\bar{\Gamma}$, but they are all linear combinations of the constraint functions. More precisely, the Einstein Hamiltonian H is of the form $H = \int_{\Sigma} (N\mathcal{H} + N^a\mathcal{H}_a)$, where N is a (positive) function on Σ , and N^a is a vector field on Σ . We then see that the observables, as defined above, are constants of the motion. The problem of observables amounts to finding enough constants of the motion to uniquely specify any vacuum spacetime (up to diffeomorphisms). Unfortunately, not a single observable is known. Indeed, aside from the work of Kuchař [7], very little is known about the form such observables can take.

Let $(q_{ab}, p^{ab}) \in \Gamma$, *i.e.*, q_{ab} is a Riemannian metric on Σ and p^{ab} is a symmetric tensor density on Σ . We will classify all *local observables*, which are defined to be constants of motion that can be expressed as integrals on Σ of densities built locally from the phase space variables (q_{ab}, p^{ab}) and their derivatives to any order. For example, the Hamiltonian H is a local observable albeit a trivial one, *i.e.*, it is equivalent to zero. Observables generate 1-parameter families of canonical transformations that preserve $\bar{\Gamma}$. Infinitesimally, the local observables generate transformations of $\bar{\Gamma}$ that are built locally from the canonical variables and their derivatives. It is not too hard to see that, in spacetime language, the existence of a local observable corresponds to the existence of an infinitesimal transformation of the *spacetime* metric that (i) is built locally from the spacetime metric and its derivatives, (ii) maps a solution of the Einstein equations to a nearby solution. Such transformations are called “generalized symmetries” by applied mathematicians. Recently all generalized symmetries of the vacuum Einstein equations have been classified [9]. They consist of a trivial scaling symmetry and the familiar diffeomorphism symmetry. The former cannot be implemented as a symplectic map of Γ , while the latter is generated by the constraint functions themselves. Because there are no other symmetries, there can be no observables (save the trivial constraints) built as local functionals of the canonical variables [10].

To summarize, we have ruled out the simplest putative resolutions of the problems of time and observables. We cannot use parametrized field theory to solve the problem of time because, strictly speaking, general relativity is not a parametrized field theory. And we have seen that there are essentially no local observables for vacuum spacetimes.

References

- [1] A. Hanson, T. Regge, C. Teitelboim, *Constrained Hamiltonian Systems*, Academia Nazionale Dei Lincei, Rome, 1976; A. Ashtekar, *Lectures on Non-Perturbative Canonical Gravity*, World Scientific, Singapore, 1991.
- [2] K. Kuchař, “Time and Interpretations of Quantum Gravity”, in the proceedings of *The Fourth Canadian Conference on General Relativity and Relativistic Astrophysics*, edited by G. Kunstatter, D. Vincent, and J. Williams, World Scientific, Singapore, 1992.
- [3] P. Bergmann, *Rev. Mod. Phys.* **33**, 510 (1961).
- [4] K. Kuchař, *J. Math. Phys.* **13**, 768 (1972).
- [5] C. Torre, *Phys. Rev. D***46**, R3231 (1992).
- [6] A. Fischer, J. Marsden, and V. Moncrief, *Ann. Inst. H. Poincaré* **33**, 147 (1980).
- [7] K. Kuchař, *J. Math. Phys.* **22**, 2640 (1981).
- [8] P. Olver, *Applications of Lie Groups to Differential Equations*, Springer-Verlag, New York, 1993.
- [9] C. Torre and I. Anderson, *Phys. Rev. Lett* **70**, 3525 (1993); *Generalized Symmetry Analysis of the Vacuum Einstein Equations*, Utah State University Preprint, 1994; *Two Component Spinors and Natural Coordinates for the Prolonged Einstein Equation Manifolds*, Utah State University Technical Report, 1994.
- [10] C. Torre, *Phys. Rev. D***48**, R2373 (1993).